

SEIZURE ACTION PLAN (SAP)



END EPILEPSY

Name: _____ Birth Date: _____
Address: _____ Phone: _____
Parent/Guardian: _____ Phone: _____
Emergency Contact/Relationship: _____ Phone: _____

Seizure Information

| Seizure Type | How Long It Lasts | How Often | What Happens |
|--------------|-------------------|-----------|--------------|
| | | | |
| | | | |
| | | | |
| | | | |

Protocol for seizure during school (check all that apply) ☒

- ☐ First aid – **Stay. Safe. Side.**
- ☐ Give rescue therapy according to SAP
- ☐ Notify parent/emergency contact
- ☐ Contact school nurse at _____
- ☐ Call 911 for transport to _____
- ☐ Other _____



First aid for any seizure

- ☐ **STAY** calm, keep calm, **begin timing seizure**
- ☐ Keep me **SAFE** – remove harmful objects, don't restrain, protect head
- ☐ **SIDE** – turn on side if not awake, keep airway clear, don't put objects in mouth
- ☐ **STAY** until recovered from seizure
- ☐ Swipe magnet for VNS
- ☐ Write down what happens _____
- ☐ Other _____

When to call 911

- ☐ Seizure with loss of consciousness longer than 5 minutes, not responding to rescue med if available
- ☐ Repeated seizures longer than 10 minutes, no recovery between them, not responding to rescue med if available
- ☐ Difficulty breathing after seizure
- ☐ Serious injury occurs or suspected, seizure in water

When to call your provider first

- ☐ Change in seizure type, number or pattern
- ☐ Person does not return to usual behavior (i.e., confused for a long period)
- ☐ First time seizure that stops on its' own
- ☐ Other medical problems or pregnancy need to be checked



When rescue therapy may be needed:

WHEN AND WHAT TO DO

If seizure (cluster, # or length) _____

Name of Med/Rx _____ How much to give (dose) _____

How to give _____

If seizure (cluster, # or length) _____

Name of Med/Rx _____ How much to give (dose) _____

How to give _____

If seizure (cluster, # or length) _____

Name of Med/Rx _____ How much to give (dose) _____

How to give _____

Care after seizure

What type of help is needed? (describe) _____

When is student able to resume usual activity? _____

Special instructions

First Responders: _____

Emergency Department: _____

Daily seizure medicine

| Medicine Name | Total Daily Amount | Amount of Tab/Liquid | How Taken (time of each dose and how much) |
|---------------|--------------------|----------------------|---|
| | | | |
| | | | |
| | | | |
| | | | |

Other information

Triggers: _____

Important Medical History _____

Allergies _____

Epilepsy Surgery (type, date, side effects) _____

Device: ☐ VNS ☐ RNS ☐ DBS Date Implanted _____

Diet Therapy ☐ Ketogenic ☐ Low Glycemic ☐ Modified Atkins ☐ Other (describe) _____

Special Instructions: _____

Health care contacts

Epilepsy Provider: _____ Phone: _____

Primary Care: _____ Phone: _____

Preferred Hospital: _____ Phone: _____

Pharmacy: _____ Phone: _____

My signature _____ Date _____

Provider signature _____ Date _____

Asthma Action Plan for Home & School

Name: _____

Birthdate: _____

Asthma Severity: ☐ Intermittent ☐ Mild Persistent ☐ Moderate Persistent ☐ Severe Persistent
☐ He/she has had many or severe asthma attacks/exacerbations

 **Green Zone** Have the child take these medicines every day, even when the child feels well.


Always use a spacer with inhalers as directed.

Controller Medicine(s): _____

Controller Medicine(s) Given in School: _____

Rescue Medicine: Albuterol/Levalbuterol _____ puffs every four hours as needed

Exercise Medicine: Albuterol/Levalbuterol _____ puffs 15 minutes before activity as needed

 **Yellow Zone** Begin the sick treatment plan if the child has a cough, wheeze, shortness of breath, or tight chest. Have the child take all of these medicines when sick.

Rescue Medicine: Albuterol/Levalbuterol _____ puffs every 4 hours as needed


Controller Medicine(s): _____

☐ Continue Green Zone medicines: _____

☐ Add: _____

☐ Change: _____

If the child is in the **yellow** zone more than **24** hours or is getting worse, follow **red** zone and call the doctor right away!

 **Red Zone** If breathing is hard and fast, ribs sticking out, trouble walking, talking, or sleeping.
Get Help Now

Take rescue medicine(s) now

Rescue Medicine: Albuterol/Levalbuterol _____ puffs every _____

Take: _____

If the child is not better right away, call 911
Please call the doctor any time the child is in the red zone.

Asthma Triggers: (List)

School Staff: Follow the Yellow and Red Zone plans for rescue medicines according to asthma symptoms.
Unless otherwise noted, the only controllers to be administered in school are those listed as "given in school" in the green zone.

- ☐ Both the asthma provider and the parent feel that the child may carry and self-administer their inhalers
☐ School nurse agrees with student self-administering the inhalers

Asthma Provider Printed Name and Contact Information:

Asthma Provider Signature:

Date:

Parent/Guardian: I give written authorization for the medications listed in the action plan to be administered in school by the nurse or other school members as appropriate. I consent to communication between the prescribing health care provider/clinic, the school nurse, the school medical advisor and school-based health clinic providers necessary for asthma management and administration of this medication.

Parent/guardian signature:

School Nurse Reviewed:

Date:

Date:

| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|

**FARE**

Food Allergy Research & Education

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: ☐ **Yes (higher risk for a severe reaction)** ☐ **No****PLACE
PICTURE
HERE****NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.****Extremely reactive to the following allergens:** _____**THEREFORE:**

- ☐ If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- ☐ If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS

**LUNG**

Shortness of breath, wheezing, repetitive cough

**HEART**

Pale or bluish skin, faintness, weak pulse, dizziness

**THROAT**

Tight or hoarse throat, trouble breathing or swallowing

**MOUTH**

Significant swelling of the tongue or lips

**SKIN**

Many hives over body, widespread redness

**GUT**

Repetitive vomiting, severe diarrhea

**OTHER**

Feeling something bad is about to happen, anxiety, confusion

**OR A
COMBINATION**
of symptoms
from different
body areas.

- 1. INJECT EPINEPHRINE IMMEDIATELY.**
 - 2. Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS**NOSE**

Itchy or runny nose, sneezing

**MOUTH**

Itchy mouth

**SKIN**

A few hives, mild itch

**GUT**

Mild nausea or discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE
SYSTEM AREA, GIVE EPINEPHRINE.**

**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: ☐ 0.1 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM

Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____



FARE
Food Allergy Research & Education

FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPi™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPi by finger grips only and slowly insert the needle into the thigh. SYMJEPi can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____